

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY) 03 Dec 2000		2. REPORT TYPE Abstract		3. DATES COVERED (From - To) 3-8 Dec 2000	
4. TITLE AND SUBTITLE Biological and hydrodynamic control of seafloor roughness: Implications to high frequency acoustic scattering				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER 062435N	
6. AUTHOR(S) Michael D. Richardson, Kevin B. Briggs, Kevin Williams, UW				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Naval Research Laboratory Marine Geosciences Division Stennis Space Center, MS 39529-5004				8. PERFORMING ORGANIZATION REPORT NUMBER  AB/7430-00-0014	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Navy Research 800 N. Quincy St. Arlington, VA 22217-5000				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT  Biological and hydrodynamic processes can both create and destroy seafloor microtopography. As part of the SAX99 experiments, natural and artificial temporal changes in seafloor roughness were monitored acoustically and quantified using bottom stereo photographs. Feeding activities of benthic megafauna and fish destroyed large-scale roughness features generated by ocean surface gravity waves within a period of weeks to months; whereas, fine-scale roughness created by raking the seafloor decayed to background levels within 24 hours. The effects of fine-scale roughness increased acoustic scattering centered at one-half the acoustic wavelength (a "Bragg" wavelength of 2 cm) by 12-18 dB in artificial manipulations of the bottom. These changes were restricted to roughness that was oriented predominantly orthogonal to the incident acoustic waves. Alternatively, seafloor roughness generated by ocean surface gravity waves had wavelengths of 50-100 cm and wave heights of 10-15 cm. These predictable large-scale roughness features should, by analogy, dramatically increase scattering at lower acoustic frequencies (near 1-2 kHz) and decay within weeks to months after storm events. [Work supported by ONR.]					
15. SUBJECT TERMS Biological, hydrodynamic, control, seafloor, roughness, SAX99, temporal changes					
16. SECURITY CLASSIFICATION OF: Abstract Unclassified			17. LIMITATION OF ABSTRACT  SAR	18. NUMBER OF PAGES  2	19a. NAME OF RESPONSIBLE PERSON Michael Richardson
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) 228-688-4621

Standard Form 298 (Rev. 8-98)  
Prescribed by ANSI Std. Z39.18

20010907 076

Biological and hydrodynamic control of seafloor roughness: implications to high-frequency acoustic scattering. Michael D. Richardson, Kevin B. Briggs (Marine Geosciences Division, Naval Research Laboratory, SSC MS 39529-5004) and Kevin L. Williams (Applied Physics Laboratory, University of Washington, 1013 NE 40<sup>th</sup> St., Seattle WA 98105-6698)

Biological and hydrodynamic processes can both create and destroy seafloor microtopography. As part of the SAX99 experiments, natural and artificial temporal changes in seafloor roughness were monitored acoustically and quantified using bottom stereo photographs. Feeding activities of benthic megafauna and fish destroyed large-scale roughness features generated by ocean surface gravity waves within a period of weeks to months; whereas, fine-scale roughness created by raking the seafloor decayed to background levels within 24 hours. The effects of fine-scale roughness increased acoustic scattering centered at one-half the acoustic wavelength (a "Bragg" wavelength of 2 cm) by 12-18 dB in artificial manipulations of the bottom. These changes were restricted to roughness that was oriented predominantly orthogonal to the incident acoustic waves. Alternatively, seafloor roughness generated by ocean surface gravity waves had wavelengths of 50-100 cm and wave heights of 10-15 cm. These predictable large-scale roughness features should, by analogy, dramatically increase scattering at lower acoustic frequencies (near 1-2 kHz) and decay within weeks to months after storm events. [Work supported by ONR.]

Number of words in abstract: 169

Suggested for special session on High-Frequency Sediment Acoustics and Associated Sediment Properties and Processes  
Technical Area: Underwater Acoustics

Special facility: Computer Projector for PowerPoint presentation  
Method of presentation: lecture

PACS Subject Classification numbers: 43.30.Hw, 43.30.Ma

Telephone Number: 228-688-4621 (M.D. Richardson) FAX: 228-688-5752

Send notice to: M.D. Richardson Email:

[mike.richardson@nrlssc.navy.mil](mailto:mike.richardson@nrlssc.navy.mil)

**NOT RELEASEABLE UNTIL**  
Security and Policy Review are  
completed and release approved  
as per SECNAVINST 5720.44A